arXiv:1402.2050v1 [math.OA] 10 Feb 2014

Finite covering projections of noncommutative torus

February 11, 2014

Petr R. Ivankov*
e-mail: * monster.ivankov@gmail.com

Abstract

This article contains is concerned with noncommutative analogue of topological finitely listed covering projections. In my previous article [4] I have already find a family of covering projections of the noncommutative torus. This article describes all covering projections of the noncommutative torus.

Contents

1 Introduction. Preliminaries **1**

2 Covering projections of noncommutative torus **4**

3 Problems **8**

4 Acknowlegment **8**

1 Introduction. Preliminaries

This article assumes elementary knowledge of following subjects

1. Algebraic topology [10].

2. C^*- algebras and operator theory [8].

Following notation is used.

Symbol	Meaning
$\mathrm{Aut}(A)$	Group * - automorphisms of C^* algebra A
A^G	Algebra of G invariants, i.e. $A^G = \{a \in A \mid ga = a, \forall g \in G\}$
$B(H)$	Algebra of bounded operators on Hilbert space H
$\mathbb{C}$ (resp. $\mathbb{R}$)	Field of complex (resp. real) numbers
$C(X)$	C^* - algebra of continuous complex valued functions on topological space X
$\mathrm{Homeo}(X)$	The space of homeomorphisms with compact-open topology [10].
$\mathcal{K}(H)$ or $\mathcal{K}$	Algebra of compact operators on Hilbert space H
$\mathrm{Map}(X, Y)$	The set of maps from X to Y
$\mathbb{T}^2$	Commutative 2-torus
$U(H) \subset \mathcal{B}(H)$	Group of unitary operators on Hilbert space H
$U(A) \in A$	Group of unitary operators of algebra A
$\mathbb{Z}$	Ring of integers
$\mathbb{Z}_m$	Ring of integers modulo m

Let us recall some definitions from my previous article [4].

1.1. *Galois extensions*. Let G be a finite group, G-Galois extensions can be regarded as particular case of Hopf-Galois extensions [6], where Hopf algebra is a commutative algebra $C(G)$. Let A be a C^*-algebra, let $G \subset \mathrm{Aut}(A)$ be a finite group of $*$- automorphisms. Let ${}_A\mathcal{M}^G$ be a category of G-equivariant modules. There is a pair of adjoint functors (F, U) given by

$$F = A \otimes_{A^G} - :_{A^G} M \to_A \mathcal{M}^G; \tag{1}$$

$$U = (-)^G :_A \mathcal{M}^G \to_{A^G} \mathcal{M}. \tag{2}$$

The unit and counit of the adjunction (F, U) are given by

$$\eta_N : N \to (A \otimes_{A^G} N)^G,\ \eta_N(n) = 1 \otimes n;$$

$$\varepsilon_M : A \otimes_{A^G} M^G \to M,\ \varepsilon_M(a \otimes m) = am.$$

Consider a map

$$\mathrm{can} : A \otimes_{A^G} A \to \mathrm{Map}(G, A) \tag{3}$$

given by

$$a_1 \otimes a_2 \mapsto (g \mapsto a_1(ga_2)),\ (a_1, a_2 \in A,\ g \in G).$$

The can is a ${}_A\mathcal{M}^G$ morphism.

Theorem 1.2. *[2] Let A be an algebra, let G be a finite group which acts on A, (F, U) functors given by (1), (2). Consider the following statements:*

1. *(F, U) is a pair of inverse equivalences;*
2. *(F, U) is a pair of inverse equivalences and $A \in_{A^G} \mathcal{M}$ is flat;*

3. *The* can *is an isomorphism and* $A \in_{A^G} \mathcal{M}$ *is faithfully flat.*

These the three conditions are equivalent.

Definition 1.3. If conditions of theorem 1.2 are hold, then A is said to be a *left faithfully flat G-Galois extension*.

Remark 1.4. Theorem 1.2 is an adapted to finite groups version of theorem from [2].

In case of commutative C^*-algebras definition 1.3 supplies algebraic formulation of finitely listed covering projections of topological spaces. However I think that above definition is not quite good analogue of noncommutative covering projections. Noncommutative algebras contains inner automorphisms. Inner automorphisms are rather gauge transformations [3] than geometrical ones. So I think that inner automorphisms should be excluded. Importance of outer automorphisms was noted by Miyashita [7]. It is reasonably take to account outer automorphisms only. I have set more strong condition.

Definition 1.5. [9] Let A be C^* - algebra. A *- automorphism α is said to be *generalized inner* if is obtained by conjugating with unitaries from multiplier algebra $M(A)$.

Definition 1.6. [9] Let A be C^* - algebra. A *- automorphism α is said to be *partly inner* if its restriction to some non-zero α- invariant two-sided ideal is generalized inner. We call automorphism *purely outer* if it is not partly inner.

Instead definitions 1.5, 1.6 following definitions are being used.

Definition 1.7. Let $\alpha \in \mathrm{Aut}(A)$ be an automorphism. A representation $\rho : A \to B(H)$ is said to be α - *invariant* if a representation ρ_α given by

$$\rho_\alpha(a) = \rho(\alpha(a)) \tag{4}$$

is unitary equivalent to ρ.

Definition 1.8. Automorphism $\alpha \in \mathrm{Aut}(A)$ is said to be *strictly outer* if for any α- invariant representation $\rho : A \to B(H)$, automorphism ρ_α is not a generalized inner automorphism.

Definition 1.9. Let A be a C^* - algebra and $G \subset \mathrm{Aut}(A)$ be a finite subgroup of * - automorphisms. An injective * - homomorphism $f : A^G \to A$ is said to be a *noncommutative finite covering projection* (or *noncommutative G - covering projection*) if f satisfies following conditions:

1. A is a finitely generated equivariant projective left and right A^G Hilbert C^*-module.
2. If $\alpha \in G$ then α is strictly outer.
3. f is a left faithfully flat G - Galois extension.

The G is said to be *covering transformation group* of f. Denote by $G(B|A)$ covering transformation group of covering projection $A \to B$.

2 Covering projections of noncommutative torus

2.1. *Noncommutative torus.* A noncommutative torus [11] A_θ is C^*-norm completion of algebra generated by two unitary elements u, v which satisfy following conditions

$$uu^* = u^*u = vv^* = v^*v = 1;$$

$$uv = e^{2\pi i\theta}vu$$

where $\theta \in \mathbb{R}$. If $\theta = 0$ then $A_\theta = A_0$ is commutative algebra of continuous functions on commutative torus $C(\mathbb{T}^2)$. There is a trace τ_0 on A_θ such that $\tau_0(\sum_{-\infty<i<\infty,-\infty<j<\infty} a_{ij}u^iv^j) = a_{00}$. C^* - norm of A_θ is defined by following way $\|a\| = \sqrt{\tau_0(a^*a)}$.

2.2. Let us recall construction from [4]. Let us consider * - homomorphism $f : A_\theta \to A_{\theta'}$, where $A_{\theta'}$ is generated by unitary elements u' and v'. Homomorphism f is defined by following way:

$$u \mapsto u'^m;$$

$$v \mapsto v'^n;$$

It is clear that

$$\theta' = \frac{\theta + k}{mn};\ (k = 0, ..., mn - 1). \tag{5}$$

Lemma 2.3. *[4] If θ is an irrational number then above $*$-homomorphism $f : A_\theta \to A_{\theta'}$ is a noncommutative covering projection.*

2.4. *Unique path lifting.* It is known that any topological covering projection $p : \tilde{X} \to X$ is a fibration with unique path lifting [10], i.e. if $\omega_1, \omega_2 : [0,1] \to \widetilde{X}$ are such that $\omega_1(0) = \omega_2(0)$ and $p(\omega_1(t)) = p(\omega_2(t))$ $(\forall t \in [0,1])$, then $\omega_1(t) = \omega_2(t)$ $(\forall t \in [0,1])$. From unique path lifting it follows that if $\alpha : [0,1] \to \mathrm{Homeo}(X)$ is a continuous map to the space of homeomorphisms such that $\alpha(0) = \mathrm{Id}_X$ then there is the unique continuous map $\tilde{\alpha} : [0,1] \to \mathrm{Homeo}(\tilde{X})$ such that $\tilde{\alpha}(0) = \mathrm{Id}_{\tilde{X}}$, and $p(\tilde{\alpha}(t)) = \alpha(t)$ $(\forall t \in [0,1])$.

Definition 2.5. Let $f : A^G \to A$ be a noncommutative covering projection. We say that f *has unique lifting* if for any continuous map $\alpha : [0,1] \to \mathrm{Aut}(A^G)$ such that $\alpha(0) = \mathrm{Id}_{A^G}$ there is a map $\tilde{\alpha} : [0,1] \to \mathrm{Aut}(A)$ such that $\tilde{\alpha}|_{A^G}(t) = \alpha(t)$ $(\forall t \in [0,1])$ and $\tilde{\alpha}(0) = \mathrm{Id}_A$.

2.6. *Action of a commutative torus.* Any point of commutative torus $\mathbb{T}^2$ can be parametrized by a pair $(z_1, z_2) \in \mathbb{C}^2$ such that $|z_1| = |z_2| = 1$. Commutative torus acts on A_θ by following way

$$u \mapsto z_1u;\ v \mapsto z_2v;\ \forall (z_1, z_2) \in \mathbb{T}^2. \tag{6}$$

Let G be a finite group and $f : A_\theta \to B$ be a G - covering projection, suppose that f has unique lifting. Then for any $(z_1, z_2) \in \mathbb{T}^2$ there is $\alpha \in \mathrm{Aut}(B)$ such that $\alpha(a) = (z_1, z_2)a$ $(\forall a \in A_\theta)$. Let $G' = \{\alpha \in \mathrm{Aut}(B) \mid \alpha|_{A_\theta} \in \mathbb{T}^2\}$. Then there is a following exact sequence of groups

$$\{e\} \to G \xrightarrow{h'} G' \xrightarrow{h} \mathbb{T}^2 \to \{e\}. \tag{7}$$

Homomorphism h is a covering projection (in topological sense) because G is a finite group. Covering projections of the commmutative torus are well known and exact sequence (7) can be rewritten by following way

$$\{e\} \to G_1 \times G_2 \xrightarrow{\mathrm{pr}_1 \times h_2'} G_1 \times G_2' \xrightarrow{h} \mathbb{T}^2 \to \{e\} \tag{8}$$

where $G = G_1 \times G_2$, $G' = G_1 \times G_2'$, G_2' is an abelian group which is isomorphic to $\mathbb{T}^2$, a homomorphism $G_1 \to \mathbb{T}^2$ $(g \mapsto h((g,e)))$ is trivial, a homomorphism $G_2' \to \mathbb{T}^2$ $(g \mapsto h((e,g)))$ is a connected covering projection of commutative torus. Sequence (8) can be decomposed into following sequences

$$\{e\} \to G_2 \xrightarrow{h_2'} G_2' \xrightarrow{h} \mathbb{T}^2 \to \{e\}; \tag{9}$$

$$\{e\} \to G_1 \to G_1 \times \mathbb{T}^2 \to \mathbb{T}^2 \to \{e\}. \tag{10}$$

Any covering of noncommutative torus can be decomposed into two covering projections which correspond to (9) and (10) respectively.
Let us consider following special cases of sequence (7):

1. (9) G' is a connected topological space $G' \to \mathbb{T}^2$ is finitely listed covering projection and G is a covering transformation group.
2. (10) $G' = G \times \mathbb{T}^2$,

2.7. *G' is a connected topological space.*
In this case $G' \approx \mathbb{T}^2$. Homomorphism $h : G' \to \mathbb{T}^2$ from (7) is given by

$$(z_1, z_2) \to (z_1^n, z_2^m)$$

where $(z_1, z_2) \in G' \approx \mathbb{T}^2$, $(n, m \in \mathbb{N})$. Any element $(z_1, z_2) \in G \subset G'$ is given by

$$(z_1, z_2) = \left(e^{\frac{2\pi i k_1}{m}} \; e^{\frac{2\pi i k_2}{n}} \right); \; (k_1, k_2 \in \mathbb{Z}).$$

Action of $G' \approx \mathbb{T}^2$ on B is an unitary representation of a compact Lie group $G' \approx \mathbb{T}^2 \to \mathrm{Aut}(B)$ [1]. Representations of $\mathbb{T}^2$ are well known, if element $b \in B$ belongs to an irreducible representation then there are $r, s \in \mathbb{Z}$ such that

$$(z_1, z_2)b = z_1^r z_1^s b; \; ((z_1, z_2) \in G' \approx \mathbb{T}^2). \tag{11}$$

An element $a \in B$ is said to be (r,s) *homogeneous* if it satisfies (11). Let $a \in B$ be a nonzero (r,s) homogeneous element, then $c = a^*a > 0$ is a $(0,0)$ homogeneous positive element, a is invariant with respect to G, i.e. $a \in B^G = A_\theta$. Any $(0,0)$ homogeneous element in A_θ is a constant, i.e. $c \in \mathbb{C}$. Moreover $c \in \mathbb{R}_+$ because c is positive element of C^*-algebra. So any (r,s) homogeneous element a satisfies following equation

$$aa^* = c; \; (\, c \in \mathbb{R}_+). \tag{12}$$

From (12) it follows that $\sqrt{c}a$ is an unitary, i.e any homogeneous element is $\mathbb{C}$ - proportional to an unitary element. Let $a_1, a_2 \in B$ be two unitary (r,s) homogeneous elements then $c = a_1 a_2^{-1}$ is a $(0,0)$ homogeneous element, i.e. $c \in \mathbb{C}$, or

$$a_1 = ca_2;\ (c \in \mathbb{C}); \tag{13}$$

From (13) that for any $(r,s) \in \mathbb{Z}^2$ a set of (r,s) homogeneous elements is a one dimensional vector space over $\mathbb{C}$. From exactness of G' action it follows that there exist a (1, 0) homogeneous nonzero element $u' \in B$. Element u'^m is G invariant and $(m,0)$ homogeneous. Element $u \in A_\theta$ is a $(m,0)$ homogeneous in B, so we have $u'^m = cu \ \ (c \in \mathbb{C})$. Similarly there is an element v' such that $v'^n = cv$. Monomials $u'^r v'^s$ are (r,s) homogeneous elements and they are $\mathbb{C}$ - generators of B. From this fact it follows that any $b \in B$ can be uniquely represented by following way

$$b = \sum_{i=0;\ j=0}^{i=m-1;\ j=n-1} a_{ij} u'^i v'^j;\ (a_{ij} \in A_\theta). \tag{14}$$

Algebra B is in fact an algebra $A_{\theta'}$ described in 2.2.

2.8. $G' = G \times \mathbb{T}^2$.
In this case we have following

$$G' \approx \bigoplus_{g \in G} \mathbb{T}^2_g$$

and a homomorphism $h : \bigoplus_{g\in G} \mathbb{T}^2_g \to \mathbb{T}^2$ is given by

$$h((t_{g_1}, ..., t_{g_n})) = t_{g_1} + ... + t_{g_n};\ (t_{g_1}, ..., t_{g_n}) \in \bigoplus_{g \in G} \mathbb{T}^2_g$$

where additive notation of the binary $\mathbb{T}^2$ group operation is used. The $\bigoplus_{g\in G} \mathbb{T}^2_g$ is a compact Lie group and any its representation is a direct sum of irreducible representations. Any irreducible representation $\bigoplus_{g\in G} \mathbb{T}^2_g \to U(\mathbb{C})$ is given by

$$((z_{1g_1}, z_{2g_1}), ..., (z_{1g_n}, z_{2g_n})) \mapsto z_{1g_1}^{i_{g_1}} z_{2g_1}^{j_{g_1}} \ ... \ z_{1g_n}^{i_{g_n}} z_{2g_n}^{j_{g_n}};$$

$$(z_{1g_k}, z_{2g_k}) \in \mathbb{T}_{g_k},\ i_{g_k}, j_{g_k} \in \mathbb{Z}.$$

An element $a \in B$ is said to be a $x = ((i_{g_1}, j_{g_1}), ..., (i_{g_n}, j_{g_n}))$ *homogeneous* if it satisfies following condition

$$((z_{1g_1}, z_{2g_1}), ..., (z_{1g_n}, z_{2g_n}))a = z_{1g_1}^{i_{g_1}} z_{2g_1}^{j_{g_1}} \ ... \ z_{1g_n}^{i_{g_n}} z_{2g_n}^{j_{g_n}} a.$$

If a' (resp. a'') is a $((i'_{g_1}, j'_{g_1}), ..., (i'_{g_n}, j'_{g_n}))$, (resp. $((i''_{g_1}, j''_{g_1}), ..., (i''_{g_n}, j''_{g_n}))$) homogeneous element then the product $a'a''$ is a $((i'_{g_1} + i''_{g_1}, j'_{g_1} + j''_{g_1}), ..., (i'_{g_n} + i''_{g_n}, j'_{g_n} + j''_{g_n}))$ homogeneous element. So B is a $\left(\mathbb{Z}^2\right)^G$ graded algebra. G naturally acts on $\left(\mathbb{Z}^2\right)^G$. If $x \in \left(\mathbb{Z}^2\right)^G$ and $a \in B$ is x - homogeneous element then ga is a gx - homogeneous element. Similarly $\mathbb{T}^2$

acts on A_θ. From this action it follows that A_θ is a $\mathbb{Z}^2$ graded algebra an element $a \in A_\theta$ is said to be (r,s) *homogeneous* if

$$(z_1, z_2)a = z_1^r z_2^s a.$$

From exactness of $\bigoplus_{g\in G} \mathbb{T}^2_g$ action it follows that there is a nonzero $((1,0),(0,0),...,(0,0))$ homogeneous element $u_{g1} \in B$. Denote by u_g a homogeneous element given by

$$u_g = g' u_{g_1},\ g' g_1 = g \in G.$$

There is the $\mathbb{C}$ - linear map $p : B \to A_\theta$ given by:

$$p(a) = \sum_{g\in G} ga;\ \forall a \in B.$$

It is clear that $p(u_{g1}) \in A_\theta$ is a $(1,0)$ homogeneous element. However any $(1,0)$ homogeneous element is equal to cu ($c \in \mathbb{C}$). If we replace u_{g_1} with $c^{-1}u_{g_1}$ then $p(u_{g_1}) = u$. From $p(u_{g1})p(u^*_{g1}) = uu^* = 1$ it follows that

$$(u_{g_1} + ... + u_{g_n})(u^*_{g_1} + ... + u^*_{g_n}) = 1. \tag{15}$$

Right part of (15) is a $((0,0),...,(0,0))$ homogeneous element in B. If $u_{g_1}u^*_{g_2} \neq 0$ then left part of (15) contains a nonzero $((1,0),(0,-1),...,(0,0))$ homogeneous summand but right part could not contain it, so we have $u_{g_1}u^*_{g_2} = 0$. Similarly we can define elements $v_{g_1},...,v_{g_n}$ and

$$v_{g1} + ... + v_{g_n} = v;$$
$$(v_{g1} + ... + v_{g_n})(v^*_{g1} + ... + v^*_{g_n}) = 1.$$

If $u_{g_1}v_{g_2} \neq 0$ then right part of

$$uv = (u_{g1} + ... + u_{g_n})(v_{g1} + ... + v_{g_n}) \tag{16}$$

contains a nonzero $((1,0),(0,1),(0,0),...,(0,0))$ homogeneous summand. However left part of (16) could not contain this summand, so we have $u_{g_1}v_{g_2} = 0$. Similarly if $g', g'' \in G$ and $g' \neq g''$ we have following:

$$u_{g'}u_{g''} = u_{g'}u^*_{g''} = u^*_{g'}u_{g''} = u^*_{g'}u^*_{g''} = v_{g'}v_{g''} = v_{g'}v^*_{g''} = v^*_{g'}v_{g''} = v^*_{g'}v^*_{g''} = 0; \tag{17}$$

$$u_{g'}v_{g''} = u_{g'}v^*_{g''} = u^*_{g'}v_{g''} = u^*_{g'}v^*_{g''} = v_{g'}u_{g''} = v_{g'}u^*_{g''} = v^*_{g'}u_{g''} = v^*_{g'}u^*_{g''} = 0. \tag{18}$$

From

$$u = u_{g_1} + ... + u_{g_n}$$

it follows that

$$u_{g_1}u^*_{g_1}u = w_1 + ... + w_n \tag{19}$$

where w_1 is a $((1,0),(0,0),...,(0,0))$ homogeneous element, w_2 is a $((0,0),(1,0),...,(0,0))$ homogeneous element, and so on. However from (17), (18) it follows that $((0,0),(1,0),...,(0,0))$,..., $((0,0),(0,0),...,(1,0))$ homogeneous summands of (19) are equal to zero, so we have

$$u_{g_1}u^*_{g_1}u = w_1 = u_{g_1}$$

or

$$e_{g_1}u = u_{g_1}$$

where $e_{g_1} = u_{g_1}u_{g_1}^*$. Similarly we can define e_g for any $g \in G$ such that

$$e_{g_1g_2} = g_1e_{g_2}. \tag{20}$$

From previous equations it follows that e_g is an idempotent for any $g \in G$ and B is a following direct sum of algebras

$$B = \bigoplus_{g \in G} e_gB.$$

A direct summand $e_gB \subset B$ is a generated by u_g, v_g subalgebra. From previous equations it follows that

$$u_gv_g = e^{2\pi i\theta}v_gu_g. \tag{21}$$

From (21) it follows that there is an isomorphism $A_\theta \to e_gB$ for any $g \in G$. In result a noncommutative covering projection f is a $*$ - homomorphism given by

$$A_\theta \to \bigoplus_{|G|} A_\theta; \tag{22}$$

$$a \mapsto (a, ..., a).$$

From (20) it follows that G just transposes direct summands of (22).

3 Problems

This construction requires condition of unique path lifting 2.5. However I do not know is this condition really necessary. Analogue of infinitely listed coverings of noncommutative torus is described in [5]. I am engaged with the general construction of infinitely listed noncommutative covering projections.

4 Acknowlegment

I would like to acknowledge a "Non-commutative geometry and topology" seminar, organized by:

1. Prof. Alexander Mishchenko,
2. Prof. Ivan Babenko,
3. Prof. Evgenij Troitsky,
4. Prof. Vladimir Manuilov,
5. Dr. Anvar Irmatov

for discussion of my work. Especially I would like acknowledge Prof. Vladimir Manuilov, because he inspired me to prove this result.

References

[1] T. Bröcker, T. Tom Dieck. *Representations of Compact Lie Groups*, Springer, Jan 1, 1985.

[2] S. Caenepeel, S. Crivei, A. Marcus, M. Takeuchi. *Morita equivalences induced by bimodules over Hopf-Galois extensions*, arXiv:math/0608572, 2007.

[3] David J. Gross. *Gauge Theory-Past, Present, and Future?* Joseph Henry Luborutoties, Ainceton University, Princeton, NJ 08544, USA. (Received November 3,1992).

[4] Petr R. Ivankov. *Noncommutative covering projections and K-homology*, arXiv:1402.0775 2014.

[5] Petr R. Ivankov. *Universal covering space of the noncommutative torus*, arXiv:1401.6748, 2014.

[6] *Lecture notes on noncommutative geometry and quantum groups*, Edited by Piotr M. Hajac.

[7] Y. Miyashita. *Finite outer Galois theory of non-commutative rings*, J. Fac. Sci. Hokkaido Univ. (I) 19 (1966), 114-134.

[8] G.J. Murhpy. *C^*-Algebras and Operator Theory.* Academic Press 1990.

[9] Marc A. Reiffel, *Actions of Finite Groups on C^* - Algebras*. Department of Mathematics University of California Berkeley. Cal. 94720 U.S.A. 1980.

[10] E.H. Spanier. *Algebraic Topology.* McGraw-Hill. New York 1966.

[11] J.C. Várilly. *An Introduction to Noncommutative Geometry*. EMS 2006.